


Available at [www.sciencedirect.com](http://www.sciencedirect.com)**ScienceDirect**journal homepage: [www.elsevier.com/locate/IJMYCO](http://www.elsevier.com/locate/IJMYCO)

# Drug development: The cell wall as a drug target

**Jonathan A.G. Cox**

School of Life &amp; Health Sciences, Aston University, Birmingham, UK

The complex and essential cell wall of *Mycobacterium tuberculosis* represents a plethora of new and old drug targets that collectively form an apparent mycobacterial “Achilles’ heel”. The mycolic acids are long-chain  $\alpha$ -alkyl- $\beta$ -hydroxy fatty acids (C<sub>70–90</sub>), which are unique to mycobacterial species, forming an integral component of the mycolyl-arabinogalactan-peptidoglycan complex. Their apparent uniqueness to the *M. tuberculosis* complex has rendered components of mycolic acid biosynthesis as powerful drug targets for specific tuberculosis (TB) chemotherapy. Here, I will discuss a contribution to TB drug discovery by deconvolution of the inhibitory mechanisms of a number of antitubercular compounds targeting mycolic acid biosynthesis. I will begin with the early days, elucidating the mode of action of ethionamide [1] and thiolactomycin [2], each targeting two separate components of the fatty acid synthase II (FAS-II) pathway. I will further discuss the recently discovered tetrahydropyrazo[1,5-*a*]pyrimidine-3-carboxamide compounds [3] which selectively target the essential, catalytically silent *M. tuberculosis* EchA6, providing a crucial lipid shunt between  $\beta$ -oxidation and FAS-II and supplying lipid precursors for essential mycolate biosynthesis. Finally, I will discuss the recent discovery of the mode of action of the indazole sulfonamides [4], inhibiting *M. tuberculosis* KasA by, a completely novel inhibitory mechanism.

## Conflict of interest

The author declared that there is no conflict of interest.

## REFERENCES

- [1] L.G. Dover et al., EthA, a common activator of thiocarbamide-containing drugs acting on different mycobacterial targets, *Antimicrob. Agents Chemother.* 51 (3) (2007) 1055–1063.
- [2] L. Kremer et al., Thiolactomycin and related analogues as novel anti-mycobacterial agents targeting KasA and KasB condensing enzymes in *Mycobacterium tuberculosis*, *J. Biol. Chem.* 275 (22) (2000) 16857–16864.
- [3] J.A. Cox et al., THPP target assignment reveals EchA6 as an essential fatty acid shuttle in mycobacteria, *Nat. Microbiol.* 1 (2016) 15006, <http://dx.doi.org/10.1038/nmicrobiol.2015.6>.
- [4] K.A. Abrahams et al., Identification of KasA as the cellular target of an anti-tubercular scaffold, *Nat. Commun.* 7 (2016) 12581, <http://dx.doi.org/10.1038/ncomms12581>.

E-mail address: [j.a.g.cox@aston.ac.uk](mailto:j.a.g.cox@aston.ac.uk)

Peer review under responsibility of Asian African Society for Mycobacteriology.  
<http://dx.doi.org/10.1016/j.ijmyco.2016.09.012>